

Effects of Luminosity Functions Induced by Relativistic Beaming on Statistics of Cosmological Gamma-Ray Bursts

Chunglee Kim, Heon-Young Chang, and Insu Yi

Korea Institute for Advanced Study, Seoul, 130-012, Korea

ciel@newton.kias.re.kr, hyc@kias.re.kr, iyi@kias.re.kr

Received _____; accepted _____

ABSTRACT

Most of gamma-ray bursts (GRBs) models have predicted that the intrinsic isotropic energy is limited to below $\sim 10^{53-54}$ ergs. Recently claimed high redshifts, correlation with supernovae, and connection to the cosmic star formation activity point to a different energy requirement and source evolution history possibly with strong beaming. We study the effects of the beaming-induced luminosity function on statistics of observed GRBs, assuming the cosmological scenario. We select and divide the BATSE 4B data into 588 long bursts ($T_{90} > 2.5$ sec) and 149 short bursts ($T_{90} < 2.5$ sec), and compare the statistics calculated in each subgroup. The $\langle V/V_{\max} \rangle$ of the long bursts is 0.2901 ± 0.0113 , and that of the short bursts is 0.4178 ± 0.0239 . For luminosity function models, we consider a cylindrical-beam and a conic-beam. We take into account the spatial distribution of GRB sources as well. A broad luminosity function is naturally produced when one introduces beaming of GRBs. We calculate the maximum detectable redshift of GRBs, $z_{\max}$. The estimated $z_{\max}$ for the cylindrical-beam case is as high as ~ 14 ($\alpha = 1.0$) and ~ 6 ($\alpha = 2.0$) for the long bursts and ~ 3 ($\alpha = 1.0$) and ~ 1.6 ($\alpha = 2.0$) for the short bursts where α is the photon index. The large $z_{\max}$ value for the short bursts is rather surprising in that the $\langle V/V_{\max} \rangle$ for this subgroup is close to the so-called Euclidean value, 0.5. We calculate the fraction of bursts whose redshifts are larger than a certain redshift z' , i.e. $f_{>z'}$. When we take $z' = 3.42$ and apply the luminosity function derived for the cylindrical-beam, the expected $f_{>z'}$ is ~ 75 % ($\alpha = 1.0$) and ~ 50 % ($\alpha = 2.0$) for long bursts. When we increase the opening angle of the conic beam to $\Delta\theta = 3^\circ.0$, $f_{>z'}$ decreases to ~ 20 % ($\alpha = 1.0$) at $z' = 3.42$. If we assume $\alpha = 2.0$, the conic-beam with $\Delta\theta = 3^\circ.0$ can not explain the redshift distribution of the observed GRBs. We conclude that the beaming-induced luminosity functions are compatible with the

redshift distribution of the observed GRBs although the apparent "Euclidean" value of $\langle V/V_{\text{max}} \rangle$ might be explained by the standard model.

Subject headings: cosmology: theory – gamma rays: bursts – methods: statistical

1. Introduction

The observation of high-redshift GRBs such as GRB970508, GRB971214, and GRB980703 (Metzger et al. 1997; Kulkarni et al. 1998; Djorgovski et al. 1998) confirmed the cosmological origin of GRBs (Paczynski 1986; Meegan et al. 1992). High redshifts ($z > 1$) of GRBs would imply a much larger apparent energy budget than the conventional energy budget $\sim 10^{53-54}$ ergs unless some significant beaming is involved (Kumar 1999; Kumar and Piran 2000). The isotropic energies of observed GRBs with known redshifts range from $\sim 10^{50}$ ergs to $\sim 10^{54}$ ergs. A broad luminosity function can be a reasonable answer for this wide range, and can be naturally expected if we assume beaming of GRBs. The suggested correlation between GRBs and certain types of supernovae (Ib/c) implies that the total energy available for GRBs could be rather limited at least in the low energy GRBs $L_{\text{int}} \sim 10^{49}$ erg/sec (Wang and Wheeler 1998; Woosley et al. 1999, see also Kippen et al. 1998; Graziani et al. 1999). If we consider the supernova explosion as a universal GRB source, then the reported GRB-supernova correlation implies that the high redshift GRBs must be strongly beamed with the beam opening solid angle, which is approximately given by

$$d\Omega \sim (L_{\text{int}}/L_{\text{obs}}) \sim 10^{-4}(L_{\text{int}}/10^{49}\text{erg/sec})(L_{\text{obs}}/10^{53}\text{erg/sec})^{-1}. \quad (1)$$

When we consider beaming, a broad luminosity range and a stringent constraint on the intrinsic energy are anticipated. Yi (1994) considered the effect of the luminosity function due to the cylindrical-beam with the inhomogeneous distribution of GRB sources

on statistics of GRBs. He analytically showed that the luminosity function follows a power law with the index $-4/3$ in the very narrow beam model when $\alpha = 1.0$. His result has been confirmed by Mao and Yi (1994) and Chang and Yi (2000). Mao and Yi (1994) studied the statistical properties of luminosity functions derived using a conic beam with a uniform spatial distribution of GRB sources. They examined the cumulative probability distribution of the peak count rates. They found that the maximum redshift increases as a product of the Lorentz factor and the opening angle decreases. We attempt to generalize the luminosity function induced by the cylindrical-beam by introducing the conical shape in such a beam. If the progenitors of GRBs were produced in the late stage of massive stars (Woosley 1993; Paczyński 1998; MacFadyen and Woosley 1999), their spatial distribution may follow the star-formation rate (SFR) of massive stars (Totani 1997; Wijers et al. 1998; Blain and Natarajan 2000). We use the recently observed SFR data to derive the possible number density distribution of GRB sources (Steidel et al. 1999).

In this paper, we study how beaming-induced luminosity functions affect the statistics of the observed GRBs in the BATSE 4B catalog (Paciesas et al. 1999) considering the two different spatial distributions of GRB sources: an SFR-motivated distribution and the uniform distribution. We employ two different SFR-motivated distributions. We discuss the effect of the photon index α ($= 1.0$ and 2.0) in the conclusion. We assume a flat universe with no cosmological constant, and adopt the Hubble constant $H_0 = 50\text{km/sec/Mpc}$ (cf. Yi 1994; Mao and Yi 1994). We analyze 775 observed bursts in the BATSE 4B catalog in order to improve the statistical significance. Assuming the bimodal distribution of durations of GRBs (Kouveliotou et al. 1993; Lamb et al. 1993; Mao, Narayan and Piran 1994; Katz and Canel 1996) we divide the total sample into two subgroups according to their duration and study the statistics of each subgroup.

The selection criteria applied to observational data are described in section 2.

Luminosity functions due to beaming are discussed in section 3. We present results and discussions in section 4. We summarise our conclusion in section 5.

2. Classification of Observational Data

We adopt the bursts in the BATSE 4B catalog (Paciesas et al. 1999) and calculate their $\langle V/V_{\max} \rangle$ for selected GRBs (Schmidt et al. 1988), using fluxes in channels 2 and 3. We choose bursts detected in 1024 ms trigger time scale with the peak count rates satisfying $C_{\max}/C_{\min} \geq 1$. Applying these criteria, we select 775 bursts among 915 bursts in the BATSE 4B $C_{\max}/C_{\min}$ table. The $\langle V/V_{\max} \rangle$ value for this total sample is 0.3177 ± 0.0102 . Then we divide the selected sample into two subgroups according to the burst durations (T_{90}), which is motivated by the reported bimodal structure in GRB durations (Kouveliotou et al. 1993; Lamb et al. 1993; Mao, Narayan and Piran 1994; Katz and Canel 1996). We define bursts with $T_{90} > 2.5$ sec as long bursts, and those with $T_{90} < 2.5$ sec as short bursts. Their estimated $\langle V/V_{\max} \rangle$ values are 0.2901 ± 0.0113 (588 bursts) and 0.4178 ± 0.0239 (149 bursts), for long bursts and short bursts respectively. The number in parentheses indicates the number of bursts belonging to each subgroup. The $\langle V/V_{\max} \rangle$ values obtained from the two subgroups show deviations from the values which are expected for a uniform distribution in a Euclidean space. In our subgroups, the long bursts have a larger deviation from the Euclidean value. It agrees with the claim by Tavani (1998). According to Tavani (1998), bursts with long durations ($T_{90} > 2.5$ sec) and hard spectra ($HR_{32} > 3.0$) show the largest deviation from the so-called Euclidean value.

3. Beaming and Luminosity Functions

We assume two kinds of number density distributions of GRB sources: (i) the SFR-motivated distribution $n_{\text{SFR}}(z)$ and (ii) the uniform distribution $n_{\text{u}}(z) = n_{\text{o,u}}$. We assume a two-sided Gaussian function as $n_{\text{SFR}}(z)$ (Kim, Yang, and Yi 1999), and fit $n_{\text{SFR}}(z)$ to the extinction-corrected SFR data (Steidel et al. 1999). The observed SFR gradually decreases or even keeps flat over a wide redshift range beyond a certain redshift (cf. Madau et al. 1998). The SFR-motivated number density distribution is approximated by

$$\begin{aligned} n_{\text{SFR}}(z) = & n_{\text{o,SFR}} \exp\left[-\frac{(z-z_c)^2}{\Delta z_1^2}\right], \quad (z < z_c) \\ & n_{\text{o,SFR}} \exp\left[-\frac{(z-z_c)^2}{\Delta z_2^2}\right], \quad (z > z_c), \end{aligned} \quad (2)$$

where Δz_1 and Δz_2 are widths of the fitting functions for the left part and the right part and $z_c = 1.5$ is a critical redshift. All parameters are determined by fitting the observational data. We set $\Delta z_1 = 1$ ($z < z_c$) and we also effectively fix $n_{\text{SFR}}(z)$ to $n_{\text{o,SFR}}$ beyond z_c . The proportional constants $n_{\text{o,SFR}}$ and $n_{\text{o,u}}$ are normalized to satisfy the number from the observational data.

3.1. Cylindrical Beaming

We consider the beaming-induced luminosity function following Yi (1993, 1994), where the relativistic beam has a constant cross-sectional area, i.e. a perfectly collimated cylindrical beam. Under this circumstance, a bulk motion of all the particles emitting the photons is parallel to the cone axis. In this case, the beam opening angle is irrelevant for the luminosity function. We assume that the beamed emission from "standard bursts" which have the same intrinsic luminosity L_{int} . The advantage of this model is that it physically defines a range of the luminosity, i.e. $[L_{\text{min}}, L_{\text{max}}]$, for a given photon index α , and the Lorentz factor γ . An apparent luminosity function $\Phi(L)$ due to the uniformly distributed

beam in space is given by

$$\Phi(L) = \frac{1}{p\beta\gamma} L_{\text{int}}^{1/p} L^{-(p+1)/p} \quad (3)$$

where $\beta = (1 - \gamma^{-2})^{1/2}$ and $p = \alpha + 2$. We adopt $\alpha = 1.0$ (Mallozzi et al. 1996) and 2.0 (Yi 1994) and the Lorentz factor $\gamma = 100$ (see Piran 1999). If we assume a bi-polar jet, the minimum and the maximum luminosities are naturally given as follows:

$$L_{\text{min}} = L_{\text{int}} B_{\text{min}}^p, \quad L_{\text{max}} = L_{\text{int}} B_{\text{max}}^p, \quad (4)$$

where $B_{\text{min}} = \gamma^{-1}$ and $B_{\text{max}} = [\gamma(1 - \beta)]^{-1}$ (Blandford and Königl 1979). The intrinsic luminosity L_{int} for each subgroup is determined by the $\langle V/V_{\text{max}} \rangle$ test. The obtained L_{int} values are $\sim 10^{43}$ erg/sec for the specific set of adopted beaming parameters. This is a much smaller value than that required in the standard candle model, which is $L_{\circ} \sim 10^{51}$ erg/sec.

We calculate two statistical quantities using the beaming-induced luminosity function explained in this subsection. First, we calculate the $\langle V/V_{\text{max}} \rangle$ as a function of threshold flux F_{th} (Che et al. 1999). The threshold flux is obtained when we put the maximum detectable redshift z_{max} in

$$F = \left(\frac{H_0^2 L}{16\pi c^2} \right) \frac{(1+z)^{1-\alpha}}{[(1+z)^{1/2} - 1]^2}, \quad (5)$$

where L is the luminosity, c is the speed of light, H_0 is the Hubble constant, and α is the photon index. After we obtain a set of F_{th} , we calculate the $\langle V/V_{\text{max}} \rangle$ for each threshold flux. The $\langle V/V_{\text{max}} \rangle$ curve converges to 0.5 as F_{th} increases sufficiently large. We convert flux to the number of bursts and plot $\langle V/V_{\text{max}} \rangle$ as a function of the number of bursts for all bursts and long and short bursts in Fig. 1 (a) for $\alpha = 1.0$ (thin lines) and $\alpha = 2.0$ (thick lines). The $\langle V/V_{\text{max}} \rangle$ vs. number of bursts relation essentially gives the same information that one obtains from the conventional $\log N - \log P$ plot. Our $\langle V/V_{\text{max}} \rangle$ test does not include the threshold effect due to the detection efficiency.

Second, we calculate the fraction of GRBs. We define the fraction of bursts located at a redshift larger than z' as

$$f_{>z'} = \frac{\text{Number of bursts with a redshift larger than } z'}{\text{Total number of bursts}}. \quad (6)$$

In Fig. 1 (b), we plot the fraction of bursts as a function of redshift for each burst subsample for the cylindrical-beam case with different number density distributions.

3.2. Conic Beaming

As a more realistic example, we adopt conically-beamed emission from GRBs, which was studied by Mao and Yi (1994), and extended by Chang and Yi (2000). We consider the luminosity function produced using three different opening angles, i.e. $\Delta\theta = 0^\circ.1, 1^\circ.0$, and $3^\circ.0$. We calculate $\langle V/V_{\text{max}} \rangle$ and the $f_{>z'}$ for each subgroup with the derived luminosity function. The bulk Lorentz factor is set to be 100. According to Mao and Yi (1994), the probability that we observe the bright bursts rapidly increases as the opening angle increases, while relatively dim bursts are not as detectable as brighter ones. Therefore, as the opening angle increases, the derived luminosity function becomes similar to that of the standard candle case, in which all bursts have the same maximum luminosity given by this luminosity function. When the opening angle is small ($\Delta\theta \ll 1/\gamma$), the luminosity function gives the same result as the cylindrical beaming case. Using this luminosity function, we calculate the same statistics as we did in previous subsection. Results are shown in Figs. 2-5 for $\alpha = 1.0$ (thin lines) and $\alpha = 2.0$ (thick lines) in the cases of each sample.

4. Results and Discussions

The left panels in Figs. 1-5 (a) show the $\langle V/V_{\text{max}} \rangle$ as a function of the number of bursts. We show results obtained from different luminosity distributions and number

density distributions. The thin lines in Fig. 1-4 represent the results obtained with $\alpha = 1.0$ and thick lines $\alpha = 2.0$. In Figs. 1-4, we assume that the SFR-motivated number density distribution, which is flat beyond a redshift larger than the critical redshift, z_c . Finally, we show our results obtained with the SFR-motivated number density distribution, which gradually decreases and $\alpha = 1.0$ in Fig. 5 (a). In all cases, we assume that $n(z)$ rapidly increases if $z < z_c$. The dotted lines represent the observed $\langle V/V_{\max} \rangle$ curves with the $\pm 3\sigma$ bound. The solid lines indicate the theoretical results derived from our luminosity functions with the SFR-motivated number density distribution of GRB sources. We plot the results obtained for the luminosity functions with the uniform distribution of burst sources as dashed lines, for comparison. Apparently all the luminosity functions we have studied satisfy the observed $\langle V/V_{\max} \rangle$ curve at the 3σ significance level. Although the number density distribution of GRB sources is believed to correlate with the SFR, we cannot rule out the uniform distribution of burst sources in this analysis (see also Krumholz et al. 1998).

The right panels in Figs. 1-5 (b) show $f_{>z'}$ in terms of z' . The y-axis corresponds to the fraction of bursts which have redshifts larger than a certain redshift z' . Solid lines result from the SFR-motivated number density distribution for the spatial distribution of GRBs and dashed lines the uniform distribution. Filled squares in Figs. 1, 3, and 5 (b) locate the known redshifts of 15 observed GRBs so far. The redshift values are quoted from http://coss.gsfc.nasa.gov/batse/counterparts/GRB_table.html. We compare the observed redshift values only with the long subsample because all GRBs with the known redshifts belong to the long bursts. It is interesting to note that, if we assume $\alpha = 1.0$, a large fraction of GRBs are distributed at high redshifts, provided that GRBs are assumed to follow the SFR, regardless of specific beaming models. However, in the case of $\alpha = 2.0$, $z_{\max}$ is significantly reduced, though it still remains quite large. The fraction for $f_{>z'}$ at $z' = 3.42$ for a cylindrical-beam case is $\sim 75\%$ for long and total samples when $\alpha = 1.0$. This is

too large a value given the fact that the observed GRBs with such high redshifts account for $\leq 10\%$ of the total bursts. This fraction become $< 50\%$ when $\alpha = 2.0$. It is also interesting to note that, though their $\langle V/V_{\max} \rangle$ value is close to a Euclidean value, most of the short bursts are distributed at high redshifts in the case of $\alpha = 1.0$. When we assume that the long bursts are uniformly distributed in space, the $z_{\max}$ estimate for the broad beaming cannot explain the largest observed redshift in both cases of $\alpha = 1.0$ and 2.0 .

Our study is based on an assumption that all GRBs have the same intrinsic luminosity and they are all beamed. Because the $\langle V/V_{\max} \rangle$ test is not sufficiently sensitive to the observed data, different luminosity functions are essentially indistinguishable. If there is an intrinsic luminosity function and the degree of beaming of GRBs is moderate, we may obtain results similar to what we have presented in this study. If all GRBs are highly beamed and their sources follow a SFR-like distribution, the maximum detectable redshift becomes very large, even in the case that the observed $\langle V/V_{\max} \rangle$ is sufficiently close to the Euclidean value, i.e., 0.5. Therefore, it is difficult to rule out a possibility that the apparent Euclidean value of $\langle V/V_{\max} \rangle$ may be due to the luminosity function, for instance, induced by beaming. In other words, the so-called Euclidean value may have nothing to do with the Euclidean distribution. If the total burst sample is a mixture of broad (or nearly non-beamed) beams and narrow beams, the exact fraction of the strongly beamed bursts may affect the burst rate estimate significantly. This is beyond the scope of this paper.

For a given Lorentz factor, the range of the derived beamed luminosity distribution becomes broader as the beam opening angle decreases. We obtain $\log L_{\max}/L_{\min} \sim 13$ in the cylindrical-beam case. However, in the case of broad beaming, $\log L_{\max}/L_{\min}$ becomes substantially smaller. This is only an effective range because the cut-off for the minimum luminosity given by the conic beam is rather loosely constrained. It is obvious that the luminosity range decreases when the opening angle increases for a fixed Lorentz factor.

This is simply explained as follows: Assuming that we observe a beamed emission with an opening angle $\Delta\theta$, if we define ψ as an angle between the line of sight and the symmetry axis of the conic beam, we can observe the beamed emission only within a certain range of ψ , $\Delta\psi$. For instance, when $1/\gamma \ll \Delta\theta$, $\Delta\psi$ is approximately $\Delta\theta$. For a given Lorentz factor, $\Delta\psi$ increases as the opening angle of cone becomes larger. The apparent luminosity range from a broad beam would decrease gradually with ψ , and in the extreme case, the isotropic radiation would have constant luminosity regardless of ψ . However, when $\Delta\theta$ is very small, the apparent luminosity rapidly decreases with ψ , and the derived luminosity range is narrower than a broad-beam case.

5. Conclusion

We have demonstrated that the beaming-induced luminosity function may explain the statistics of the observed GRBs. In the case of the cylindrically beamed luminosity function, we determined the intrinsic luminosity L_{int} by the $\langle V/V_{\text{max}} \rangle$ test for each subsample. In all cases, the intrinsic luminosity is much smaller than that of non-beamed luminosity functions, i.e. $L \sim 10^{52-53}$ erg/sec. The maximum luminosity obtained from our luminosity functions, in both the cylindrical-beam and the conic-beam cases, is $L_{\text{max}} \sim 10^{50-51}$ erg/sec. This is compatible with the required value for isotropic radiation. The ratio between the maximum luminosity for the long bursts and that for the short bursts $L_{\text{max,long}}/L_{\text{max,short}}$ is ~ 7 for the narrow beam and ~ 3 for the broad beam, respectively. This implies that these two subgroups may have two different intrinsic luminosities, and hence probably two different origin. The obtained z_{max} in the cylindrical-beam case are $\sim 10, 14, 3$ ($\alpha = 1.0$) and $\sim 4.5, 5.5, 1.6$ ($\alpha = 2.0$) for the total selected sample, and long and short subgroup, respectively. The obtained z_{max} in the broader beam case (i.e. $\Delta\theta = 3^\circ.0$) are $\sim 4, 6, 2$ ($\alpha = 1.0$) and $\sim 2.2, 2.6, 1.2$ ($\alpha = 2.0$) for the total selected sample, and long and short

subgroup, respectively. The maximum known redshift we consider, $z = 3.42$, is estimated for GRB971214. However, even a larger value of $z < 3.9$ is suggested for GRB980329 (http://coss.gsfc.nasa.gov/batse/counterparts/GRB_table.html). Without any detailed information on the redshift distribution of GRBs, the maximum redshift gives a rather strong constraint on the ratio between the Lorentz factor and the opening angle $\Delta\theta$. For the conic beam case, the luminosity function derived for the narrow opening angle seems to fit the $\langle V/V_{\text{max}} \rangle$ curve better.

From this simple analysis we have carried out, we have shown that the beaming-induced luminosity function may account for the basic statistical properties of the observed GRBs. The apparent Euclidean distribution of GRBs may be an indication of the presence of the GRB luminosity function.

Although it is beyond the scope of this work, it is interesting to point out that there could be a potentially observable correlation between spectral hardness and luminosity since the most Doppler-boosted burst would probe the hardest part of the spectrum. Any wide scatters in intrinsic spectral shapes would make the detection of this correlation complicated.

We thank H. Kim and K. Kwak for useful discussions. IY is supported in part by the KRF grant No. 1998-001-D00365. We appreciate the referee, Ralph A. M. J. Wijers, for his suggestions and some useful information.

REFERENCES

- Blain, A.W. & Natarajan, P. 2000, MNRAS, 312, 35
- Blandford, R. D. & Königl, A. 1979, ApJ, 232, 34
- Chang, H.-Y. & Yi, I. 2000, astro-ph/0005302
- Che, H., Yang, Y. & Nemiroff, R. J. 1999, ApJ, 516, 559
- Djorgovski, S. G. et al. 1998, ApJ, 508, L17
- Graziani, C., Lamb, D. Q., and Marion, G. H. 1999, A&AS, 138, 469
- Katz, J. I. & Canel, L. M. 1996, ApJ, 471, 915
- Kim, C., Yang, J., and Yi, I. 1999, J. of Korean Phys. Soc., 34, 459
- Kippen, R. M. et al. 1998, ApJ, 504, L27
- Kouveliotou, C. et al. 1993, ApJ, 413, L101
- Krumholz, M. et al. 1998, ApJ, 506, L81
- Kulkarni, S. R. et al. 1998, Nature, 393, 35
- Kumar, P. 1999, ApJ, 523, L113
- Kumar, P. and Piran, T. 2000, ApJ, 535, 152
- Lamb, D. Q., Graziani, C., and Smith, I. A. 1993, ApJ, 413, L11
- MacFadyen, A. I. & Woosley, S. E. 1999, ApJ, 524, 262
- Madau, P., Pozzetti, L., and Dickinson, M. 1998, ApJ, 498, 10
- Mallozzi, R. S., Pendleton, G. N., and Paciesas, W. S. 1996, ApJ, 471, 636

- Mao, S., Narayan, R. and Piran, T. 1994, ApJ, 420, 171
- Mao, S., & Yi, I. 1994, ApJ, 424, L131
- Meegan, C. A., et al. 1992, Nature, 355, 143
- Metzger, M. R. et al. 1997, Nature, 387, 878
- Paciesas, W. S. et al. 1999, 1999, ApJS, 122, 465
- Paczynski, B. 1986, ApJ, 308, L51
- Paczynski, B. 1998, ApJ, 494, L45
- Piran, T. 1999, Phys. Rep., 314, 575
- Schmidt, M., Higdon, J. C., and Hueter, G. 1988, ApJ, 329, L85
- Steidel, C., Adelberger, K. L., Giavalisco, M., and Dickinson, M. 1999, ApJ, 519, 1
- Tavani, M. 1998, ApJ, 497, 915
- Totani, T. 1997, ApJ, 486, L71
- Wang, L. & Wheeler, J. C. 1998, ApJ, 504, L87
- Wijers, R. A. M. J. 1998, MNRAS, 294, L13
- Woosley, S. E. 1993, ApJ, 405, 273
- Woosley, S. E. et al. 1999, ApJ, 516, 788
- Yi, I. 1993, Phys. Rev. D, 48, 4518
- Yi, I. 1994, ApJ, 431, 543

Fig. 1.— Two statistical properties of three sample sets for the cylindrical-beam case are shown. Data sets are indicated at the upper right corners in each panel. In the left panels, we plot the $\langle V/V_{\max} \rangle$ curve as a function of number of bursts. Dotted lines represent the observed $\langle V/V_{\max} \rangle$ with $\pm 3\sigma$ bounds. Solid lines and dashed lines show theoretical results, the former result from the SFR-motivated number density distribution for the spatial distribution of GRBs and the latter the uniform distribution. In the right panels, we show the $f_{>z'}$ in terms of z' for each sample set. The y-axis is the fraction of GRBs which would be located farther than a certain redshift z' . Solid lines and dashed lines are same as in the left panels. Filled squares indicate the estimated redshifts of 15 observed GRBs. In all panels, the photon index α is assumed to be 1.0 (thin lines) and 2.0 (thick lines). Note that the $z_{\max}$ is significantly reduced when $\alpha = 2.0$.

Fig. 2.— Plots similar to Fig. 1. Results for the total sample for different opening angles in the conic beam model. Opening angles are indicated at the upper right corners in each panel. In all panels, the photon index α is assumed to be 1.0 (thin lines) and 2.0 (thick lines).

Fig. 3.— Same plots as Fig. 2. for the long bursts. In all panels, the photon index α is assumed to be 1.0 (thin lines) and 2.0 (thick lines).

Fig. 4.— Same plots as Fig. 2 for the short bursts. In all panels, the photon index α is assumed to be 1.0 (thin lines) and 2.0 (thick lines).

Fig. 5.— Same plots as Fig. 2 for the conic-beam case for the total sample. We adopt a bi-Gaussian function as a number density function of GRB sources, $n(z)$, which is different from that used in Figs. 1-4. It gradually decreases at high redshifts. The photon index α is assumed to be 1.0.









